

FROM FACTORIZATIONS OF NONCOMMUTATIVE POLYNOMIALS TO COMBINATORIAL TOPOLOGY

VLADIMIR RETAKH

To D.B. Fuchs on the occasion of his 70-th anniversary

ABSTRACT. This is an extended version of a talk given by the author at the conference “Algebra and Topology in Interaction” on the occasion of the 70th Anniversary of D.B. Fuchs at UC Davis in September 2009. It is a brief survey of an area originated around 1995 by I. Gelfand and the speaker.

0. INTRODUCTION

Factorization of polynomials is one of the most fundamental and most classical problems in mathematics. We know a lot about factorizations of polynomials over different fields and, more generally, over commutative rings. Much less is known about factorizations of polynomials over noncommutative rings, e.g. polynomials with matrix coefficients. Unlike their commutative counterparts, noncommutative polynomials admit many different factorizations. This makes their theory much harder and more interesting.

In 1995 I. Gelfand and the speaker constructed $n!$ different factorizations of a noncommutative polynomial in one variable with n roots in “generic” position. Later with R. Wilson we studied “algebras of pseudo-roots” or “noncommutative splitting algebras” associated with such factorizations. These algebras can be described in terms of special directed graphs (quivers) called layered graphs.

To any cell complex one can also associate a layered graphs and a “splitting algebra” defined by this graph. There are surprising connections between properties of cell complexes and related splitting algebras. Here I construct a bridge between noncommutative algebra related to factorizations of polynomials and combinatorial topology.

Key words and phrases. noncommutative polynomials, directed graphs, Koszul algebras, Hilbert series, combinatorial topology.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

I. FACTORIZATIONS OF NONCOMMUTATIVE POLYNOMIALS

1.1. Let us start with a secondary school problem. You are given a square polynomial

$$x^2 + a_1x + a_2$$

over an associative (noncommutative) unital ring and its roots x_1, x_2 . How do you generalize the Vieté formulas $-a_1 = x_1 + x_2$ and $a_2 = x_1x_2$?

Let us assume that the difference $x_1 - x_2$ is invertible and set

$$x_{1,2} = (x_2 - x_1)x_2(x_1 - x_2)^{-1}, \quad x_{2,1} = (x_1 - x_2)x_1(x_2 - x_1)^{-1}.$$

Proposition 1.1.

$$-a_1 = x_{1,2} + x_1 = x_{2,1} + x_2$$

and

$$a_2 = x_{1,2}x_1 = x_{2,1}x_2$$

(note the order of multipliers in the last formula!)




FIGURE 1

Relations between $x_{1,2}, x_{2,1}, x_1, x_2$ can be described by the following graph (see Fig. 1). Here sums and products corresponding to the different paths from vertex $\{12\}$ to $\emptyset$ coincide. One can also present these relations in a unified way by setting $e_1 = x_{1,2}$, $e_2 = x_1$, $f_1 = x_{2,1}$, $f_2 = x_2$:

$$(t - x_{1,2})(t - x_1) = (t - x_{2,1})(t - x_2)$$

where t is a central variable.

1.2. We proceed now to polynomials of any degree over a unital associative ring. Denote by $[1, n]$ the set of indices $\{1, 2, \dots, n\}$. Assume that the polynomial

$$x^n + a_1 x^{n-1} + \dots + a_{n-1} x + a_0$$

has n roots $x_1, x_2, \dots, x_n$ in a *generic* position, i.e. for any subset $\{i_1, i_2, \dots, i_{k+1}\}$ of $[1, n]$, $k \geq 1$, the Vandermonde matrix

$$W(i_1, i_2, \dots, i_{k+1}) = \begin{pmatrix} x_{i_1}^k & x_{i_2}^k & \dots & x_{i_{k+1}}^k \\ x_{i_1}^{k-1} & x_{i_2}^{k-1} & \dots & x_{i_{k+1}}^{k-1} \\ \dots & \dots & \dots & \dots \\ 1 & 1 & \dots & 1 \end{pmatrix}$$

is invertible.

For the matrix $W(i_1, i_2, \dots, i_{k+1})$ denote the first row without $x_{i_{k+1}}^k$ by r and the last column without $x_{i_{k+1}}^k$ by c . Set

$$w_{i_1 i_2 \dots i_{k+1}} = x_{i_{k+1}}^k - r \cdot W(i_1, i_2, \dots, i_k)^{-1} \cdot c.$$

The element $w_{i_1 i_2 \dots i_{k+1}}$ is one of the *quasideterminants* of the matrix $W(i_1, i_2, \dots, i_{k+1})$ (see [GR1-5, GGRW, RW1, Os1-2]).

Example 1.2. One can see that $w_{i_1 i_2} = x_{i_1} - x_{i_2}$.

The element $w_{i_1 i_2 \dots i_{k+1}}$ does not depend on the ordering of the first k indices $i_1, i_2, \dots, i_k$, therefore we denote it by $w_{A_k, i_{k+1}}$ where $A_k = \{i_1, i_2, \dots, i_k\}$.

Assume now that the element $w_{A_k, i_{k+1}}$ is invertible and set

$$x_{A_k, i_{k+1}} = w_{A_k, i_{k+1}} \cdot x_{i_{k+1}} \cdot w_{A_k, i_{k+1}}^{-1}.$$

We are now ready to generalize Proposition 1.1.

Theorem 1.3. ([GR4-5]) *Let $i_1, i_2, \dots, i_n$ be an ordering of $1, 2, \dots, n$. Under our assumptions*

$$\begin{aligned} -a_1 &= \sum_{k=1}^n x_{A_{k-1}, i_k}, \\ (1.1) \quad a_2 &= \sum_{k > \ell} x_{A_{k-1}, i_k} x_{A_{\ell-1}, i_\ell}, \\ &\dots \\ (-1)^n a_n &= x_{A_{n-1}, i_n} x_{A_{n-2}, i_{n-1}} \dots x_{A_0, i_1}. \end{aligned}$$

Here $A_0 = \emptyset$ and $x_{A_0, i_1} = x_{i_1}$.

Remark. In the commutative case the right-hand sides of these formulas are elementary symmetric functions in $x_1, x_2, \dots, x_n$. In the noncommutative case the right-hand sides are also symmetric in these variables (but you have to keep the order of multipliers!). This remark is a cornerstone of the theory of noncommutative symmetric functions developed in [GKLLRT] and subsequent papers.

The formulas of Theorem 1.3 can be unified in the following way. Let t be a central variable. Set $P(t) = t^n + a_1 t^{n-1} + \dots + a_{n-1} t + a_0$. Then

$$(1.2) \quad P(t) = (t - x_{A_{n-1}, i_n})(t - x_{A_{n-2}, i_{n-1}}) \dots (t - x_{A_0, i_1})$$

and in a generic case there are $n!$ factorizations of $P(t)$ defined by different orderings of $1, 2, \dots, n$.

From now on we consider polynomials $P(t)$ over a ring R . An element $x \in R$ is a *left* root of $P(t)$ if $P(t)$ is divisible by $t - x$ from the left, and a *right* root of $P(t)$ if $P(t)$ is divisible by $t - x$ from the right. In this terminology every $x_{\emptyset, i}$ is a left root and every $x_{12 \dots \hat{i} \dots n, i}$ is a right root of $P(t)$.

Generally speaking, we cannot interpret elements $x_{A, i}$ as roots, and so we call them *pseudo-roots* due to the following definition (see [RSW3]).

Definition 1.4. An element $x \in R$ is a *pseudo-root* of a polynomial $P(t)$ if there exist polynomials $S(t), T(t) \in R[t]$ such that

$$P(t) = S(t)(t - x)T(t).$$

1.3. We construct now a universal algebra of pseudo-roots Q_n defined by generators and relations (see [GRW1]). The algebra Q_n is generated by elements $x_{A, i}$ where $A \subset [1, n]$ and $i \in [1, n] \setminus A$. The relations in Q_n are defined by the triples (A, i, j) where $A \subset [1, n]$ and $i, j \in [1, n] \setminus A$, $i \neq j$:

$$(1.3a) \quad x_{A \cup \{i\}, j} + x_{A, i} = x_{A \cup \{j\}, i} + x_{A, j},$$

$$(1.3b) \quad x_{A \cup \{i\}, j} \cdot x_{A, i} = x_{A \cup \{j\}, i} + \cdot_{A, j}.$$

Relations (1.3) can be united by the equality

$$(1.4) \quad (t - x_{A \cup \{i\}, j})(t - x_{A, i}) = (t - x_{A \cup \{j\}, i})(t - x_{A, j})$$

where t is a formal central parameter.

Equality (1.4) implies that the products (1.2) where $A_k = \{i_1, i_2, \dots, i_k\}$ do not depend on the orderings of $\{1, 2, \dots, n\}$ and, therefore, define the monic polynomial $P(t)$ over Q_n . The generators $x_{A,i}$ are the pseudo-roots of $P(t)$.

Note that relation (1.3) can be described geometrically by using the diamond graph from Fig 1 where the sets $\{1, 2\}$, $\{1\}$, $\{2\}$, $\emptyset$ are replaced by $A \cup \{i, j\}$, $A \cup \{i\}$, $A \cup \{j\}$, and A respectfully.

The next theorem shows that algebra Q_n is a large algebra containing at least $n! + 1$ free subalgebras with n generators. Any ordered set $I = \{i_1, i_2, \dots, i_n\}$ defines a factorization (1.2) with pseudo-roots x_{A_{k-1}, i_k} , $k = 1, 2, \dots, n$. Denote by Q_I the subalgebra generated by these pseudo-roots.

Theorem 1.4. [GRW1] *For each I algebra Q_I is a free subalgebra of Q_n . The intersection Sym_n of free subalgebras $\cap Q_I$ over all orderings of $\{1, 2, \dots, n\}$ is again the free subalgebra generated by elements $a_1, a_2, \dots, a_n$ defined by formulas (1.1).*

One may consider elements $(-1)^k a_k$ as noncommutative analogs of the commutative elementary functions and algebra Sym_n as an analogue of the corresponding commutative symmetric algebra. As we already said, a theory of noncommutative symmetric algebras was developed in a series of papers starting with [GKLLRT].

1.4. To confirm that Q_n has a large growth, we will pick a set of linearly independent generators parametrized by subsets $B = (j_1, j_2, \dots, j_k) \subset [1, n]$. The generators are

$$x_B = x_{\emptyset, j_1} + x_{j_1, j_2} + \dots + x_{j_1, j_2, \dots, j_{k-1}, j_k}.$$

Linear relations (1.3a) imply that x_B does not depend on the order of indices $j_1, \dots, j_k$. Elements x_B generate algebra Q_n and satisfy only quadratic relations (see ([GRW1])). Therefore, Q_n is a quadratic algebra with a natural grading defined by $\deg x_B = 1$. Let d_i be the dimension of the i -th component of Q_n according to this grading. Then according to [GRW1, GGRSW] the graded dimension or the Hilbert series $H(Q_n, \tau) = \sum_{i \geq 0} d^i \tau^i$ of the algebra Q_n is

$$(1.5) \quad H(Q_n, \tau) = \frac{1 - \tau}{1 - \tau(2 - \tau)^n}.$$

2. FROM GRAPHS TO NONCOMMUTATIVE POLYNOMIALS

2.1. In the previous section we used graphs to describe relations between pseudo-roots. Now we will use graphs to construct polynomials with noncommutative coefficients and their pseudo-roots (see [GRSW, GGRW, RSW3] and other papers).

Let $\Gamma = (V, E)$ be a directed graph (quiver) where V is the set of vertices and E is the set of edges. For any $e \in E$ denote by $t(e)$ the tail of e and by $h(e)$ the head of e .

Suppose that Γ is a *layered* graph, i.e. $V = \coprod_{i=0}^n V_i$ and $t(e) \in V_i$, $i > 0$, implies $h(e) \in V_{i-1}$. We call n the *height* of the graph and we write $|v| = i$ if $v \in V_i$. Set $V_+ = \coprod_{i>0} V_i$.

Examples 2.1.

a) Let V be the set of subsets of the set $[1, n]$. For any $v \in V$ set $|v|$ to be the cardinality of v . An edge e goes from v to w if w is the subset of v such that $|w| = |v| - 1$. In this case $V_0 = \{\emptyset\}$ and $V_n = \{[1, n]\}$.

b) Let V be the set of linear subspaces of a vector space F of dimension n . For any $v \in V$ set $|v|$ to be the dimension of v . An edge e goes from v to w if w is the subspace of v such that $|w| = |v| - 1$. In this case V_0 consists of the zero subspace and $V_n = \{F\}$.

c) This is a generalization of Example 2.1a. Let X be a regular cell complex. Construct the corresponding layered graph $\Gamma_X = (V, E)$ as follows. Let V be the set of cells $\sigma \in X$. Set $|\sigma| = \dim \sigma + 1$. An edge $e \in E$ goes from σ to τ if and only if τ lies in the closure of σ and $|\tau| = |\sigma| - 1$. In this case $V_0 = \emptyset$ as in Example 1.

2.2. Following [GRSW] we construct now an algebra $A(\Gamma)$ associated with a layered graph Γ . A path π in Γ is a sequence of edges $\pi = (e_1, e_2, \dots, e_k)$ such that $h(e_i) = t(e_{i+1})$ for $i = 1, 2, \dots, k-1$. We call $t(e_1)$ the tail of π and denote it by $t(\pi)$ and we call $h(e_k)$ the head of π and denote it by $h(\pi)$. The algebra $A(\Gamma)$ is generated over a given field by generators $e \in E$ subject to the following relations. Let t be a formal parameter commuting with edges $e \in E$. Any two paths $\pi = (e_1, e_2, \dots, e_k)$ and $\pi' = (f_1, f_2, \dots, f_k)$ with the same tail and head define the relation

$$(2.1) \quad (t - e_1)(t - e_2) \dots (t - e_k) = (t - f_1)(t - f_2) \dots (t - f_k).$$

In fact, relation (2.1) is equivalent to k relations

$$e_1 + e_2 + \dots + e_k = f_1 + f_2 + \dots + f_k,$$

$$(2.2) \quad \begin{aligned} \sum_{i < j} e_i e_j &= \sum_{i < j} f_i f_j, \\ &\dots \\ e_1 e_2 \dots e_k &= f_1 f_2 \dots f_k. \end{aligned}$$

We call $A(\Gamma)$ the *splitting* algebra associated with graph Γ . The terminology is justified by the following considerations. Assume that there are only one vertex $*$ of the minimal level 0 and only one vertex M of the maximal level n , and that for any edge e there exists a path $\theta = (e_1, e_2, \dots, e_n)$ from M to $*$ containig e . Set

$$P(t) = (t - e_1)(t - e_2) \dots (t - e_n).$$

Then $P(t)$ is a polynomial over $A(\Gamma)$, any edge in the graph Γ corresponds to a pseudo-root of $P(t)$, and any path from the maximal to minimal vertex corresponds to a factorization of $P(t)$ into a product of linear factors.

Example 2.2. Let Γ be the graph defined in Example 2.1a. Then $A(\Gamma)$ is the algebra Q_n defined in Section 1.3.

2.3. We will discuss now some properties of algebras $A(\Gamma)$. First we describe the Hilbert series of such algebras assuming that the layered graph Γ has exactly one minimal vertex $*$ and for any vertex $v \in V_i, i > 0$ there is a path $\pi_v = (h_1, h_2, \dots, h_i)$ from v to the minimal vertex. The images of new elements $r_v = h_1 + h_2 + \dots + h_i$ in $A(\Gamma)$ do not depend on the choice of path from v to $*$ and are linearly independent. According to formulas (2.2) elements r_v generate algebra $A(\Gamma)$ and satisfy homogeneous relations of order two and higher. By assigning the degree one to each element r_v we obtain a natural grading on $A(\Gamma)$. The corresponding Hilbert series $H(A(\Gamma), \tau)$ is well defined, provided that the set of vertices V is finite.

To write an expression for the Hilbert series we introduce a *graded Möbius function* on Γ . We define a partial order on the set of vertices of Γ by putting $v > w$ if and only if there is a path from v to w . The classical Möbius function μ on $V \times V$ can be defined in the following way: $\mu(w, v) = 0$ if $w \not\leq v$, $\mu(v, v) = 1$, and if $w < v$

$$\mu(w, v) = \sum_{w=v_0 < v_1 < \dots < v_\ell = v} (-1)^\ell.$$

We introduce the *graded Möbius function* of the graph as series

$$\mathcal{M}(\Gamma, \tau) = \sum_{w < v} \mu(v, w) \tau^{|v| - |w|}.$$

The following theorem was proved in [RSW2].

Theorem 2.3. *Let $\Gamma = (V, E)$ be a layered graph with a unique minimal element $*$ of level 0. Assume that the set of vertices is finite. Then*

$$H(\Gamma, \tau) = \frac{1 - \tau}{1 - \tau \mathcal{M}(\Gamma, \tau)}.$$

One can show (see [RSW4]) that the inverse series $H(\Gamma, \tau)^{-1}$ is, in fact, a polynomial. The degree of the polynomial equals to the height of Γ . Theorem 2.3 implies formula (1.5) for algebra Q_n .

Generalized layered graphs corresponding to factorizations of polynomials into products of not necessary linear factors were studied in [RW2].

3. QUADRATIC SPLITTING ALGEBRAS $A(\Gamma)$ AND THEIR DUAL ALGEBRAS

3.1. Let $\Gamma = (V, E)$ be a layered graph. As in Subsection 2.3 we assume that the layered graph Γ has exactly one minimal vertex and that for any vertex $v \in V_i$, $i > 0$ there is a path $\pi_v = (h_1, h_2, \dots, h_i)$ from v to the minimal vertex. In this case the splitting algebra $A(\Gamma)$ is defined by a set of homogeneous relations of order 2 and higher. Following [RSW1] we describe now a condition when algebra $A(\Gamma)$ is defined by quadratic relations only.

We say that vertices v, v' of the same level $i > 0$ are connected by a *down-up* sequence if there exist vertices $v = v_0, v_1, v_2, \dots, v_k = v' \in V_i$ and $w_1, w_2, \dots, w_k \in V_{i-1}$ such that $w_j < v_{j-1}, v_j$ for $j = 1, 2, \dots, k$. According to [SRW1], the layered graph Γ is *uniform* if for any pair of edges $e, e' \in E$ with a common tail, $t(e) = t(e')$, their heads $h(e), h(e')$ are connected by a down-up sequence. It was proved in [RSW1] that if the graph Γ is uniform then the splitting algebra $A(\Gamma)$ is quadratic, i.e. defined by relations of order 2.

3.2. For a quadratic algebra A over a field F there is a notion of the dual quadratic algebra $A^!$. To define $A^!$ denote by W the vector space of generators of A and by $R \subset W \otimes W$ the linear space of relations of A . Denote by W^* the dual space of W and by $R^\perp$ the annihilator of R in $W^* \otimes W^*$. Algebra $A^!$ is a quadratic algebra defined by generators from W^* and relations from $R^\perp$. It is well-known (see, for example, [PP]) that an algebra A is Koszul if and only if its dual algebra $A^!$ is. In this case their Hilbert series are connected by the simple equation

$$(3.1) \quad H(A, \tau)H(A^!, -\tau) = 1.$$

We call a quadratic algebra A *numerically Koszul* (see [RSW4]) if the Hilbert series of algebras A and $A^!$ satisfy equation (3.1).

3.3. Assuming that a layered graph Γ is uniform one can describe the dual algebra $A(\Gamma)^!$ in terms of vertices and edges of the graph (see [RSW3]). We describe now a slightly different algebra $B(\Gamma)$.

There is a natural filtration on $A(\Gamma)$ defined by the ranking function $|\cdot|$. The corresponding graded algebra is also quadratic. Its dual algebra $B(\Gamma)$ can be described in the following way (see [CPS]). For any $v \in V_+$ let $S(v)$ be the set of all vertices $w \in V$ such that there is an edge going from v to w .

Theorem 3.1. *The algebra $B(\Gamma)$ is generated by vertices $v \in V_+$ subject to the relations:*
i) $u \cdot v = 0$ if there is no edge going from u to v ;
ii) $v \cdot \sum_{w, w \in S(v)} w = 0$.

If the set of vertices V is finite, the algebra $B(\Gamma)$ is finite-dimensional. Algebras $B(\Gamma)$ were studied in [CPS]. According to the general theory $A(\Gamma)$ is Koszul if and only if $B(\Gamma)$ is Koszul. If these algebras are Koszul then $H(B(\Gamma), \tau) = H(A(\Gamma), -\tau)^{-1}$.

Since not all splitting algebras are Koszul, we are going to investigate the difference between $H(B(\Gamma), \tau)$ and $H(A(\Gamma), -\tau)^{-1}$ for layered graphs Γ .

Let $\Gamma = (V, E)$ be a layered graph, $V = \coprod_{i=0}^n V_i$. Denote by the same letter Γ the partially ordered set (poset) associated to the graph, by $\Delta(\Gamma)$ the *order* complex of the poset $[\Gamma]$, and by $H^i(\Gamma, F)$ the cohomology groups of $\Delta(\Gamma)$ with the coefficients in a field F . Let $b_i(\Gamma)$ be the dimension of $H^i(\Gamma, F)$.

For a vertex $v \in V$ and $k \leq |v|$ denote by $\Gamma_{v,k}$ the graph induced by all vertices $w < v$, $|w| \geq |v| - k + 1$ with the added minimal vertex $*$, $|*| = 0$.

Note that $H(B(\Gamma), \tau)$ is a polynomial in τ of degree n .

The following theorem was obtained in [RSW5].

Theorem 3.2. *The coefficient of τ^k in $H(A(\Gamma), -\tau)^{-1} - H(B(\Gamma), \tau)$ equals*

$$\sum_{v \in V_\ell, \ell \geq k} \sum_{j=0}^{\ell-1} b_{\ell-j-1}(\Gamma_{v,k}).$$

4. SPLITTING ALGEBRAS ASSOCIATED TO CELL COMPLEXES

4.1. Splitting algebras associated to cell complexes provide many of examples of Koszul and non-Koszul algebras. The study of such algebras was originated in [RSW4] and continued in [CPS, SS].

Recall that to any regular cell complex X one can associate the layered graph Γ_X (see Example 2.1c).

Theorem 4.1. *Let X be a finite cell complex. Algebras $A(\Gamma_X)$ and $B(\Gamma_X)$ defined over any ground field are Koszul.*

This theorem was proved in [RSW4] for the abstract simplicial complexes and in [CPS] for any CW -complexes. The Koszulity of the algebra Q_n was earlier proved in [SW, Pi]. Our construction gives a large number of Koszul algebras of exponential growth as well as finite-dimensional Koszul algebras.

4.2. Assume now that X is a finite cell complex of dimension n . We say that X is *pure* if X is the closure of its cells of dimension n . Following [CPS] we say that a pure complex X is *connected by codimension one faces* if any pair of cells of dimension n is connected by a down-up sequence.

Let X be a pure cell complex of dimension n connected by codimension one faces. By adding to the graph Γ_X one maximal vertex with the obvious rank we obtain the graph $\hat{\Gamma}_X$.

$A(\Gamma_X)$ and $A(\hat{\Gamma}_X)$ have completely different properties. The following theorem was proved in [RSW4] for two-dimensional manifolds and later in [CPS] in general case.

Theorem 4.2. *Let X be a pure and regular CW -complex of dimension n that is connected by codimension one faces. Let F be a field.*

- i) If algebras $A(\hat{\Gamma}_X)$ and $B(\hat{\Gamma}_X)$ over the ground field F are Koszul, then $H^k(X, F) = 0$ for all $0 < k < n$.
- ii) If X is a manifold and $H^k(X, F) = 0$ for all $0 < k < n$ then algebras $A(\hat{\Gamma}_X)$ and $B(\hat{\Gamma}_X)$ are Koszul over F .

Theorem 4.2 implies that the Koszul property of algebras $A(\hat{\Gamma}_X)$ and $B(\hat{\Gamma}_X)$ unlike the algebras $A(\Gamma_X)$ and $B(\Gamma_X)$ over the ground field depends on the field. As pointed in [CPS], if X is a regular cellular representation of $\mathbb{RP}^2$, then $A(\hat{\Gamma}_X)$ and $B(\hat{\Gamma}_X)$ are Koszul if the field F does not have characteristic 2.

Remark. Note that Theorem 3.2. can be used to describe the discrepancy between Hilbert series of algebras $A(\hat{\Gamma}_X, -\tau)^{-1}$ and $B(\hat{\Gamma}_X, \tau)$ via Betti numbers of the cell complex X . It shows that splitting algebras $A(\hat{\Gamma}_X)$ and $B(\hat{\Gamma}_X)$ carry a lot of information about topological properties of the cell complex X . Note also that Theorem 3.2 can be used to prove a weaker version of Theorem 4.2.

4.3. Theorem 4.2 gives a combinatorial solution to the question when algebras $A(\hat{\Gamma}_X)$ and $B(\hat{\Gamma}_X)$ are Koszul. These algebras are combinatorial invariants but not topological invariants. In a recent paper [SS] it was shown, however, that the property of these algebras to be Koszul is a topological invariant. The following theorem was proved in [SS].

Theorem 4.3. *Let X be a pure regular cell complex of dimension n connected through codimension one faces. Then algebras $A(\hat{\Gamma}_X)$ and $B(\hat{\Gamma}_X)$ are Koszul for a field F if and only if both of the following conditions hold:*

- i) $\tilde{H}_i(X; F) = 0$ for $i < n$;
- ii) $\tilde{H}_i(X, X - \{p\}; F) = 0$ for each $p \in X$ and each $i < n$.

Theorem 4.3 shows that the Koszul property for algebras $A(\hat{\Gamma}_X)$ and $B(\hat{\Gamma}_X)$ is a homeomorphic invariant. There are, however, homotopy equivalent, pure regular cell complexes X and Y such that $A(\hat{\Gamma}_X)$ is Koszul but $A(\hat{\Gamma}_Y)$ is not (see [SS]).

REFERENCES

- [B] A. Björner, *Topological Methods*, Handbook of Combinatorics, 1995, pp. 1819–1872.
- [CPS] T. Cassidy, C. Phan, B. Shelton, *Noncommutative Koszul algebras from combinatorial topology*, arXiv:0811.3450, J. Reine Angew. Math. (to appear).
- [GGRSW] I. Gelfand, S. Gelfand, V. Retakh, S. Serconek, and R. Wilson, *Hilbert series of quadratic algebras associated with decompositions of noncommutative polynomials*, J. Algebra **254** (2002), 279–299.
- [GGRW1] I. Gelfand, S. Gelfand, V. Retakh, R. Wilson, *Quasideterminants*, Advances in Math. **193** (2005), no. 1, 56–141.
- [GGRW2] I. Gelfand, S. Gelfand, V. Retakh, R. Wilson, *Factorizations of polynomials over noncommutative algebras and sufficient sets of edges in directed graphs*, Lett. Math. Physics **74** (2005), no. 2, 153–167.

- [GKLLRT] I. Gelfand, D. Krob, A. Lascoux, B. Leclerc, V. Retakh, and J.-Y. Thibon, *Noncommutative symmetric functions*, Advances in Math. **112** (1995), no. 2, 218–348.
- [GR1] I. Gelfand and V. Retakh, *Determinants of matrices over noncommutative rings*, Funct. Anal. Appl. **25** (1991), no. 2, 91–102.
- [GR2] ———, *A theory of noncommutative determinants and characteristic functions of graphs*, Funct. Anal. Appl. **26** (1992), no. 4, 1–20.
- [GR3] ———, *A theory of noncommutative determinants and characteristic functions of graphs. I*, Publ. LACIM, UQAM, Montreal **14** (1993), 1–26.
- [GR4] ———, *Noncommutative Vieta theorem and symmetric functions*, Gelfand Mathematical Seminars 1993–95, Birkhäuser, Boston, 1996, pp. 93–100.
- [GR5] ———, *Quasideterminants*, I, Selecta Math. (N.S.) **3** (1997), no. 4, 517–546.
- [GRSW] I. Gelfand, V. Retakh, S. Serconek, R. Wilson, *On a class of algebras associated to directed graphs*, Selecta Math. **11** (2005), no. 2, 281–295.
- [GRW1] I. Gelfand, V. Retakh, and R. Wilson, *Quadratic-linear algebras associated with decompositions of noncommutative polynomials and differential polynomials*, Selecta Math. (N.S.) **7** (2001), 493–523.
- [Os1] B. L. Osofsky, *Quasideterminants and right roots of polynomials over division rings*, Algebras, rings and their representations, World Sci. Publ., 2006, pp. 241–263.
- [Os2] ———, *Noncommutative linear algebra*, Algebra and its Applications, Contemp. Math. **419** (2006), AMS, 231–254.
- [Pi] D. Piontkovski, *Algebras associated to pseudo-roots of noncommutative polynomials are Koszul*, Intern. J. Algebra Comput., **15** (2005), 643–648.
- [PP] A. Polishchuk, L. Positselski, *Quadratic algebras*, Amer. Math. Society, Providence, RI, 2005.
- [RSW1] V. Retakh, S. Serconek, R. Wilson, *On a class of Koszul algebras associated to directed graphs*, J. of Algebra **304** (2006), 1114–1129.
- [RSW2] ———, *Hilbert series of algebras associated to directed graphs*, J. of Algebra **312** (2007), 142–151.
- [RSW3] ———, *Construction of some algebras associated to directed graphs and related to factorizations of noncommutative polynomials*, Proceedings of the Conference "Lie Algebras, Vertex Operator Algebras and Their Applications", Contemp. Math **442** (2007).
- [RSW4] ———, *Koszulity of splitting algebras associated with cell complexes*, arXiv:0810.1241, J. of Algebra (to appear).
- [RSW5] ———, *Hilbert series of some algebras associated with cell complexes and cohomology* (to appear).
- [RW1] V. Retakh and R. Wilson, *Advanced course on quasideterminants and universal localization*. 124 pp., CRM, Barcelona, 2007.
- [RW2] ———, *Algebras associated to directed acyclic graphs*, Advances in Appl. Math **42** (2009), 42–59.
- [SS] H. Sadofsky and B. Shelton, *The Koszul property as a topological invariant and measure of singularities*, arXiv:0911.2541.
- [SW] S. Serconek and R. L. Wilson, *Quadratic algebras associated with decompositions of noncommutative polynomials are Koszul algebras*, J. Algebra **278** (2004), 473–493.

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, PISCATAWAY, NJ 08854
E-mail address: vretakh@math.rutgers.edu